

EXPRESS MAIL NO. EV354970770US

**APPLICATION
FOR
UNITED STATES LETTERS PATENT**

Applicants: Samuel Achilefu, Raghavan Rajagopalan,
Richard B. Dorshow, Joseph E. Bugaj,
Hermo N. Jimenez, and Muthunadar P. Periasamy

Title: LIGHT SENSITIVE COMPOUNDS FOR INSTANT
DETERMINATION OF ORGAN FUNCTION

Assignee: Mallinckrodt Inc.

Beverly A. Lyman, Esq.
WOOD, HERRON & EVANS, L.L.P.
2700 Carew Tower
441 Vine Street
Cincinnati, OH 45202-2917
(513) 241-2324-Office
(513) 421-7269-Facsimile

Attorney Docket No.: MRD/57CP

SPECIFICATION

**LIGHT SENSITIVE COMPOUNDS FOR
INSTANT DETERMINATION OF ORGAN FUNCTION**

This application is a Continuation-In-Part of U.S. Patent Application Serial No. 09/688,942 filed October 16, 2000, now pending and expressly incorporated by reference herein in its entirety.

FIELD OF THE INVENTION

This invention relates to novel optical probes for use in physiological function monitoring, particularly indole and benzoindole compounds.

5 **BACKGROUND OF THE INVENTION**

Dynamic monitoring of physiological functions of patients at the bedside is highly desirable in order to minimize the risk of acute renal failure brought about by various clinical, physiological, and pathological conditions (C.A. Rabito, L.S.T. Fang, and A.C. Waltman, Renal function in patients at risk

10 with contrast material-induced acute renal failure: Noninvasive real-time

monitoring, *Radiology* 1993, 186, 851-854; N.L. Tilney, and J.M. Lazarus, Acute renal failure in surgical patients: Causes, clinical patterns, and care, *Surgical Clinics of North America*, 1983, 63, 357-377; B.E. VanZee, W.E. Hoy, and J.R. Jaenike, Renal injury associated with intravenous pyelography in non-
5 diabetic and diabetic patients, *Annals of Internal Medicine*, 1978, 89, 51- 54; S. Lundqvist, G. Edbom, S. Groth, U. Stendahl, and S. O. Hietala, Iohexol
clearance for renal function measurement in gynecologic cancer patients, *Acta*
Radiologica, 1996, 37, 582-586; P. Guesry, L. Kaufman, S. Orlof, J.A. Nelson, S. Swann, and M. Holliday, Measurement of glomerular filtration rate by
10 fluorescent excitation of non-radioactive meglumine iothalamate, *Clinical*
Nephrology, 1975, 3, 134-138). This monitoring is particularly important in the case of critically ill or injured patients because a large percentage of these patients face the risk of multiple organ failure (MOF), resulting in death (C.C. Baker et al., Epidemiology of Trauma Deaths, *American Journal of Surgery*,
15 1980, 144-150; G. Regel et al., Treatment Results of Patients with Multiple Trauma: An Analysis of 3406 Cases Treated Between 1972 and 1991 at a
German Level I Trauma Center, *Journal of Trauma*, 1995, 38, 70-77). MOF is a sequential failure of lung, liver, and kidneys, and is incited by one or more severe causes such as acute lung injury (ALI), adult respiratory distress
20 syndrome (ARDS), hypermetabolism, hypotension, persistent inflammatory focus, or sepsis syndrome. The common histological features of hypotension and shock leading to MOF include tissue necrosis, vascular congestion, interstitial and cellular edema, hemorrhage, and microthrombi. These changes affect the lung, liver, kidneys, intestine, adrenal glands, brain, and pancreas, in
25 descending order of frequency (J. Coalson, *Pathology of Sepsis, Septic Shock*,

and Multiple Organ Failure, New Horizons: Multiple Organ Failure Syndrome, D.J. Bihari and F.B. Cerra (Eds.), *Society of Critical Care Medicine*, Fullerton, CA, 1986, pp. 27-59). The transition from early stages of trauma to clinical MOF is marked by the extent of liver and renal failure and a change in mortality risk from about 30% to about 50% (F.B. Cerra, Multiple Organ Failure Syndrome. New Horizons: Multiple Organ Failure, D.J. Bihari and F.B. Cerra (Eds). *Society of Critical Care Medicine*, Fullerton, CA, 1989, pp. 1-24).

Serum creatinine measured at frequent intervals by clinical laboratories is currently the most common way of assessing renal function and following the dynamic changes in renal function which occur in critically ill patients (P.D. Doolan, E.L. Alpen, and G.B. Theil, A clinical appraisal of the plasma concentration and endogenous clearance of creatinine, *American Journal of Medicine*, 1962, 32, 65-79; J.B. Henry (Ed). Clinical Diagnosis and Management by Laboratory Methods, 17th Edition, W.B. Saunders, Philadelphia, PA, 1984); C.E. Speicher, The right test: A physician's guide to laboratory medicine, W.B. Saunders, Philadelphia, PA, 1989). These values are frequently misleading, since age, state of hydration, renal perfusion, muscle mass, dietary intake, and many other clinical and anthropometric variables affect the value. In addition, a single value returned several hours after sampling is difficult to correlate with other important physiologic events such as blood pressure, cardiac output, state of hydration and other specific clinical events (e.g., hemorrhage, bacteremia, ventilator settings and others). An approximation of glomerular filtration rate can be made via a 24-hour urine collection, but this requires 24 hours to collect the sample, several more hours to analyze the sample, and a meticulous bedside collection technique. New or

repeat data are equally cumbersome to obtain. Occasionally, changes in serum creatinine must be further adjusted based on the values for urinary electrolytes, osmolality, and derived calculations such as the "renal failure index" or the "fractional excretion of sodium." These require additional samples
5 of serum collected contemporaneously with urine samples and, after a delay, precise calculations. Frequently, dosing of medication is adjusted for renal function and thus can be equally as inaccurate, equally delayed, and as difficult to reassess as the values upon which they are based. Finally, clinical decisions in the critically ill population are often as important in their timing as they are in
10 their accuracy.

Exogenous markers such as inulin, iohexol, ^{51}Cr -EDTA, Gd-DTPA, or $^{99\text{m}}\text{Tc}$ -DTPA have been reported to measure the glomerular filtration rate (GFR) (P.L. Choyke, H.A. Austin, and J.A. Frank, Hydrated clearance of gadolinium-DTPA as a measurement of glomerular filtration rate, *Kidney International*, 1992, 41, 1595-1598; M.F. Tweedle, X. Zhang, M. Fernandez, P. Wedeking, A.D. Nunn, and H.W. Strauss, A noninvasive method for monitoring renal status at bedside, *Invest. Radiol.*, 1997, 32, 802-805; N. Lewis, R. Kerr, and C. Van Buren, Comparative evaluation of urographic contrast media, inulin, and $^{99\text{m}}\text{Tc}$ -DTPA clearance methods for determination of glomerular filtration
15 rate in clinical transplantation, *Transplantation*, 1989, 48, 790-796). Other markers such as ^{123}I and ^{125}I labeled o-iodohippurate or $^{99\text{m}}\text{Tc}$ -MAG₃ are used to assess tubular secretion process (W.N. Tauxe, Tubular Function, in *Nuclear Medicine in Clinical Urology and Nephrology*, W.N. Tauxe and E.V. Dubovsky, Editors, pp. 77-105, Appleton Century Crofts, East Norwalk, 1985; R. Muller-Suur, and C. Muller-Suur, Glomerular filtration and tubular secretion of MAG₃ in
20
25

rat kidney, *Journal of Nuclear Medicine*, 1989, 30, 1986-1991). However, these markers have several undesirable properties such as the use of radioactivity or *ex-vivo* handling of blood and urine samples. Thus, in order to assess the status and to follow the progress of renal disease, there is a considerable
5 interest in developing a simple, safe, accurate, and continuous method for determining renal function, preferably by non-radioactive procedures. Other organs and physiological functions that would benefit from real-time monitoring include the heart, the liver, and blood perfusion, especially in organ transplant patients.

10 Hydrophilic, anionic substances are generally recognized to be excreted by the kidneys (F. Roch-Ramel, K. Besseghir, and H. Murer, Renal excretion and tubular transport of organic anions and cations, *Handbook of Physiology, Section 8, Neurological Physiology, Vol. II*, E.E. Windhager, Editor, pp. 2189-2262, Oxford University Press, New York, 1992; D.L. Nosco, and J.A.
15 Beaty-Nosco, Chemistry of technetium radiopharmaceuticals 1: Chemistry behind the development of technetium-^{99m} compounds to determine kidney function, *Coordination Chemistry Reviews*, 1999, 184, 91-123). It is further recognized that drugs bearing sulfonate residues exhibit improved clearance through the kidneys (J. Baldas, J. Bonnyman, Preparation, HPLC studies and
20 biological behavior of technetium-^{99m} and ^{99m}TcN0-radiopharmaceuticals based on quinoline type ligands, *Nucl. Med. Biol.*, 1999, 19, 491-496; L. Hansen, A. Taylor, L., L.G. Marzilli, Synthesis of the sulfonate and phosphonate derivatives of mercaptoacetyl triglycine. X-ray crystal structure of
Na₂[ReO(mercaptoacetyl glycyglycyl aminomethane-sulfonate)]·3H₂O, *Met.-*
25 *Based Drugs*, 1994, 1, 31-39).

Assessment of renal function by continuously monitoring the blood clearance of exogenous optical markers, viz., fluorescein bioconjugates derived from anionic polypeptides, has been developed by us and by others (R.B. Dorshow, J.E. Bugaj, B.D. Burleigh, J.R. Duncan, M.A. Johnson, and W.B. Jones, Noninvasive fluorescence detection of hepatic and renal function, *Journal of Biomedical Optics*, 1998, 3, 340-345; M. Sohtell et al., FITC-Inulin as a Kidney Tubule Marker in the Rat, *Acta. Physiol. Scand.*, 1983, 119, 313-316, each of which is expressly incorporated herein by reference). The main drawback of high molecular weight polypeptides is that they are immunogenic. In addition, large polymers with narrow molecular weight distribution are difficult to prepare, especially in large quantities. Thus, there is a need in the art to develop low molecular weight compounds that absorb and/or emit light that can be used for assessing renal, hepatic, cardiac and other organ functions.

SUMMARY OF THE INVENTION


The present invention overcomes these difficulties by incorporating hydrophilic anionic or polyhydroxy residues in the form of sulfates, sulfonates, sulfamates, phosphates, polyethers, polyamino carboxylic acids and their derivatives, and strategically positioned hydroxyl groups. Thus, the present invention is related to novel dyes containing multiple hydrophilic moieties and their use as diagnostic agents for assessing organ function and functional status of tumors.

The novel compounds of the present invention comprise dyes of Formulas 1 to 6 which are hydrophilic and absorb light in the visible and near infrared regions of the electromagnetic spectrum. The blood clearance rate can be modified by formulating the dyes in liposomes, micelles, or other

microparticles. This enhances their use for physiological monitoring of many organs. The ease of modifying the clearance pathways of the dyes after *in vivo* administration permits their use for physiological monitoring. Compounds with longer blood persistence are useful for angiography and organ perfusion analysis, which is particularly useful in organ transplant and critical ill patients. Predominant kidney clearance of the dyes enables their use for dynamic renal function monitoring, and rapid liver uptake of the dyes from blood serves as a useful index for the evaluation of hepatic function.

As illustrated in Figures 1-7, these dyes are designed to inhibit aggregation in solution by preventing intramolecular and intermolecular induced hydrophobic interactions.

The present invention relates particularly to the novel compounds comprising indoles of the general Formula 1




Formula 1

wherein R₃, R₄, R₅, R₆, and R₇, and Y₁ are independently selected from the group consisting of -H, C1-C10 alkoxyl, C1-C10 polyalkoxyalkyl, C1-C20 polyhydroxyalkyl, C5-C20 polyhydroxyaryl, glucose derivatives of R groups, saccharides, amino, C1-C10 aminoalkyl, cyano, nitro, halogen, hydrophilic peptides, arylpolysulfonates, C1-C10 alkyl, C1-C10 aryl, -SO₃T, -CO₂T, -OH, -(CH₂)_aSO₃T, -(CH₂)_aOSO₃T, -(CH₂)_aNHSO₃T, -(CH₂)_aCO₂(CH₂)_bSO₃T,

- $-(CH_2)_aOCO(CH_2)_bSO_3T$, $-(CH_2)_aCONH(CH_2)_bSO_3T$,
 $-(CH_2)_aNHCO(CH_2)_bSO_3T$, $-(CH_2)_aNHCONH(CH_2)_bSO_3T$,
 $-(CH_2)_aNHCSNH(CH_2)_bSO_3T$, $-(CH_2)_aOCONH(CH_2)_bSO_3T$, $-(CH_2)_aPO_3HT$,
 $-(CH_2)_aPO_3T_2$, $-(CH_2)_aOPO_3HT$, $-(CH_2)_aOPO_3T_2$, $-(CH_2)_aNHPO_3HT$,
5 $-(CH_2)_aNHPO_3T_2$, $-(CH_2)_aCO_2(CH_2)_bPO_3HT$, $-(CH_2)_aCO_2(CH_2)_bPO_3T_2$,
 $-(CH_2)_aOCO(CH_2)_bPO_3HT$, $-(CH_2)_aOCO(CH_2)_bPO_3T_2$,
 $-(CH_2)_aCONH(CH_2)_bPO_3HT$, $-(CH_2)_aCONH(CH_2)_bPO_3T_2$,
 $-(CH_2)_aNHCO(CH_2)_bPO_3HT$, $-(CH_2)_aNHCO(CH_2)_bPO_3T_2$,
 $-(CH_2)_aNHCONH(CH_2)_bPO_3HT$, $-(CH_2)_aNHCONH(CH_2)_bPO_3T_2$,
10 $-(CH_2)_aNHCSNH(CH_2)_bPO_3HT$, $-(CH_2)_aNHCSNH(CH_2)_bPO_3T_2$,
 $-(CH_2)_aOCONH(CH_2)_bPO_3HT$, and $-(CH_2)_aOCONH(CH_2)_bPO_3T_2$, $-CH_2(CH_2-O$
 $-CH_2)_c-CH_2-OH$, $-(CH_2)_d-CO_2T$, $-CH_2-(CH_2-O-CH_2)_e-CH_2-CO_2T$, $-(CH_2)_f-NH_2$,
 $-CH_2-(CH_2-O-CH_2)_g-CH_2-NH_2$, $-(CH_2)_h-N(R_a)-(CH_2)_i-CO_2T$, and $-(CH_2)_j-N(R_b)$
 $-CH_2-(CH_2-O-CH_2)_k-CH_2-CO_2T$; W_1 is selected from the group consisting of
15 $-CR_cR_d$, $-O-$, $-NR_c$, $-S-$, and $-Se$; a , b , d , f , h , i , and j independently vary from 1-
10; c , e , g , and k independently vary from 1-100; R_a , R_b , R_c , and R_d are defined
in the same manner as Y_1 ; T is either H or a negative charge.

The present invention also relates to the novel compounds comprising benzoindoles of general Formula 2




Formula 2

- wherein R_8 , R_9 , R_{10} , R_{11} , R_{12} , R_{13} , R_{14} , and Y_2 are independently selected from the group consisting of -H, C1-C10 alkoxy, C1-C10 polyalkoxyalkyl, C1-C20 polyhydroxyalkyl, C5-C20 polyhydroxyaryl, glucose derivatives of R groups,
- 5 saccharides, amino, C1-C10 aminoalkyl, cyano, nitro, halogen, hydrophilic peptides, arylpolysulfonates, C1-C10 alkyl, C1-C10 aryl, $-\text{SO}_3\text{T}$, $-\text{CO}_2\text{T}$, $-\text{OH}$, $-(\text{CH}_2)_a\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{OSO}_3\text{T}$, $-(\text{CH}_2)_a\text{NHSO}_3\text{T}$, $(\text{CH}_2)_a\text{CO}_2(\text{CH}_2)_b\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{OCO}(\text{CH}_2)_b\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{CONH}(\text{CH}_2)_b\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{NHCO}(\text{CH}_2)_b\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{NHCONH}(\text{CH}_2)_b\text{SO}_3\text{T}$,
- 10 $-(\text{CH}_2)_a\text{NHCSNH}(\text{CH}_2)_b\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{OCONH}(\text{CH}_2)_b\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{PO}_3\text{HT}$, $-(\text{CH}_2)_a\text{PO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{OPO}_3\text{HT}$, $-(\text{CH}_2)_a\text{OPO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{NHPO}_3\text{HT}$, $-(\text{CH}_2)_a\text{NHPO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{CO}_2(\text{CH}_2)_b\text{PO}_3\text{HT}$, $-(\text{CH}_2)_a\text{CO}_2(\text{CH}_2)_b\text{PO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{OCO}(\text{CH}_2)_b\text{PO}_3\text{HT}$, $-(\text{CH}_2)_a\text{OCO}(\text{CH}_2)_b\text{PO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{CONH}(\text{CH}_2)_b\text{PO}_3\text{HT}$, $-(\text{CH}_2)_a\text{CONH}(\text{CH}_2)_b\text{PO}_3\text{T}_2$,
- 15 $-(\text{CH}_2)_a\text{NHCO}(\text{CH}_2)_b\text{PO}_3\text{HT}$, $-(\text{CH}_2)_a\text{NHCO}(\text{CH}_2)_b\text{PO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{NHCONH}(\text{CH}_2)_b\text{PO}_3\text{HT}$, $-(\text{CH}_2)_a\text{NHCONH}(\text{CH}_2)_b\text{PO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{NHCSNH}(\text{CH}_2)_b\text{PO}_3\text{HT}$, $-(\text{CH}_2)_a\text{NHCSNH}(\text{CH}_2)_b\text{PO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{OCONH}(\text{CH}_2)_b\text{PO}_3\text{HT}$, and $-(\text{CH}_2)_a\text{OCONH}(\text{CH}_2)_b\text{PO}_3\text{T}_2$, $-\text{CH}_2(\text{CH}_2-\text{O}$

-CH₂)_c-CH₂-OH, -(CH₂)_d-CO₂T, -CH₂-(CH₂-O-CH₂)_e-CH₂-CO₂T, -(CH₂)_f-NH₂,
 -CH₂-(CH₂-O-CH₂)_g-CH₂-NH₂, -(CH₂)_h-N(R_a)-(CH₂)_i-CO₂T, and -(CH₂)_j-N(R_b)
 -CH₂-(CH₂-O-CH₂)_k-CH₂-CO₂T; W₂ is selected from the group consisting of
 -CR_cR_d, -O-, -NR_c, -S-, and -Se; a, b, d, f, h, i, and j independently vary from 1-
 5 10; c, e, g, and k independently vary from 1-100; R_a, R_b, R_c, and R_d are defined
 in the same manner as Y₂; T is either H or a negative charge.

The present invention also relates to the novel compounds
 comprising cyanine dyes of general Formula 3




Formula 3

10 wherein R₁₅, R₁₆, R₁₇, R₁₈, R₁₉, R₂₀, R₂₁, R₂₂, R₂₃, Y₃, and Z₃ are independently
 selected from the group consisting of -H, C1-C10 alkoxy, C1-C10
 polyalkoxyalkyl, C1-C20 polyhydroxyalkyl, C5-C20 polyhydroxyaryl, glucose
 derivatives of R groups, saccharides, amino, C1-C10 aminoalkyl, cyano, nitro,
 halogen, hydrophilic peptides, arylpolysulfonates, C1-C10 alkyl, C1-C10 aryl,
 15 -SO₃T, -CO₂T, -OH, -(CH₂)_aSO₃T, -(CH₂)_aOSO₃T,
 -(CH₂)_aNHSO₃T, -(CH₂)_aCO₂(CH₂)_bSO₃T, -(CH₂)_aOCO(CH₂)_bSO₃T,
 -(CH₂)_aCONH(CH₂)_bSO₃T, -(CH₂)_aNHCO(CH₂)_bSO₃T,
 -(CH₂)_aNHCONH(CH₂)_bSO₃T, -(CH₂)_aNHCSNH(CH₂)_bSO₃T,
 -(CH₂)_aOCONH(CH₂)_bSO₃T, -(CH₂)_aPO₃HT, -(CH₂)_aPO₃T₂, -(CH₂)_aOPO₃HT,

- $-(CH_2)_aOPO_3T_2$, $-(CH_2)_aNHPO_3HT$, $-(CH_2)_aNHPO_3T_2$,
 $-(CH_2)_aCO_2(CH_2)_bPO_3HT$, $-(CH_2)_aCO_2(CH_2)_bPO_3T_2$, $-(CH_2)_aOCO(CH_2)_bPO_3HT$,
 $-(CH_2)_aOCO(CH_2)_bPO_3T_2$, $-(CH_2)_aCONH(CH_2)_bPO_3HT$,
 $-(CH_2)_aCONH(CH_2)_bPO_3T_2$, $-(CH_2)_aNHCO(CH_2)_bPO_3HT$,
5 $-(CH_2)_aNHCO(CH_2)_bPO_3T_2$, $-(CH_2)_aNHCONH(CH_2)_bPO_3HT$,
 $-(CH_2)_aNHCONH(CH_2)_bPO_3T_2$, $-(CH_2)_aNHCSNH(CH_2)_bPO_3HT$,
 $-(CH_2)_aNHCSNH(CH_2)_bPO_3T_2$, $-(CH_2)_aOCONH(CH_2)_bPO_3HT$, and
 $-(CH_2)_aOCONH(CH_2)_bPO_3T_2$, $-CH_2(CH_2-O-CH_2)_c-CH_2-OH$, $-(CH_2)_d-CO_2T$,
 $-CH_2-(CH_2-O-CH_2)_e-CH_2-CO_2T$, $-(CH_2)_f-NH_2$, $-CH_2-(CH_2-O-CH_2)_g-CH_2-NH_2$,
10 $-(CH_2)_h-N(R_a)-(CH_2)_i-CO_2T$, and $-(CH_2)_j-N(R_b)-CH_2-(CH_2-O-CH_2)_k-CH_2-CO_2T$;
 W_3 and X_3 are selected from the group consisting of $-CR_cR_d$, $-O-$, $-NR_c$, $-S-$,
and $-Se$; V_3 is a single bond or is selected from the group consisting of $-O-$,
 $-S-$, $-Se-$, and $-NR_a$; a , b , d , f , h , i , and j independently vary from 1-10; c , e , g ,
and k independently vary from 1-100; a_3 and b_3 vary from 0 to 5; R_a , R_b , R_c , and
15 R_d are defined in the same manner as Y_3 ; T is either H or a negative charge.

The present invention further relates to the novel compounds comprising cyanine dyes of general Formula 4




Formula 4

- wherein R_{24} , R_{25} , R_{26} , R_{27} , R_{28} , R_{29} , R_{30} , R_{31} , R_{32} , R_{33} , R_{34} , R_{35} , R_{36} , Y_4 , and Z_4 are independently selected from the group consisting of -H, C1-C10 alkoxy, C1-C10 polyalkoxyalkyl, C1-C20 polyhydroxyalkyl, C5-C20 polyhydroxyaryl, glucose derivatives of R groups, saccharides, amino, C1-C10 aminoalkyl, cyano, nitro, halogen, hydrophilic peptides, arylpolysulfonates, C1-C10 alkyl, C1-C10 aryl, $-\text{SO}_3\text{T}$, $-\text{CO}_2\text{T}$, $-\text{OH}$, $-(\text{CH}_2)_a\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{OSO}_3\text{T}$, $-(\text{CH}_2)_a\text{NHSO}_3\text{T}$, $-(\text{CH}_2)_a\text{CO}_2(\text{CH}_2)_b\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{OCO}(\text{CH}_2)_b\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{CONH}(\text{CH}_2)_b\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{NHCO}(\text{CH}_2)_b\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{NHCONH}(\text{CH}_2)_b\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{NHCSNH}(\text{CH}_2)_b\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{OCONH}(\text{CH}_2)_b\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{PO}_3\text{HT}$, $-(\text{CH}_2)_a\text{PO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{OPO}_3\text{HT}$, $-(\text{CH}_2)_a\text{OPO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{NHPO}_3\text{HT}$, $-(\text{CH}_2)_a\text{NHPO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{CO}_2(\text{CH}_2)_b\text{PO}_3\text{HT}$, $-(\text{CH}_2)_a\text{CO}_2(\text{CH}_2)_b\text{PO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{OCO}(\text{CH}_2)_b\text{PO}_3\text{HT}$, $-(\text{CH}_2)_a\text{OCO}(\text{CH}_2)_b\text{PO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{CONH}(\text{CH}_2)_b\text{PO}_3\text{HT}$, $-(\text{CH}_2)_a\text{CONH}(\text{CH}_2)_b\text{PO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{NHCO}(\text{CH}_2)_b\text{PO}_3\text{HT}$, $-(\text{CH}_2)_a\text{NHCO}(\text{CH}_2)_b\text{PO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{NHCONH}(\text{CH}_2)_b\text{PO}_3\text{HT}$, $-(\text{CH}_2)_a\text{NHCONH}(\text{CH}_2)_b\text{PO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{NHCSNH}(\text{CH}_2)_b\text{PO}_3\text{HT}$, $-(\text{CH}_2)_a\text{NHCSNH}(\text{CH}_2)_b\text{PO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{OCONH}(\text{CH}_2)_b\text{PO}_3\text{HT}$, and $-(\text{CH}_2)_a\text{OCONH}(\text{CH}_2)_b\text{PO}_3\text{T}_2$, $-\text{CH}_2(\text{CH}_2\text{-O-CH}_2)_c\text{-CH}_2\text{-OH}$, $-(\text{CH}_2)_d\text{-CO}_2\text{T}$, $-\text{CH}_2\text{-(CH}_2\text{-O-CH}_2)_e\text{-CH}_2\text{-CO}_2\text{T}$, $-(\text{CH}_2)_f\text{-NH}_2$, $-\text{CH}_2\text{-(CH}_2\text{-O-CH}_2)_g\text{-CH}_2\text{-NH}_2$, $-(\text{CH}_2)_h\text{-N(R}_a\text{)-(CH}_2)_i\text{-CO}_2\text{T}$, and $-(\text{CH}_2)_j\text{-N(R}_b\text{)-CH}_2\text{-(CH}_2\text{-O-CH}_2)_k\text{-CH}_2\text{-CO}_2\text{T}$;
- W_4 and X_4 are selected from the group consisting of $-\text{CR}_c\text{R}_d$, $-\text{O-}$, $-\text{NR}_c$, $-\text{S-}$, and $-\text{Se}$; V_4 is a single bond or is selected from the group consisting of $-\text{O-}$, $-\text{S-}$, $-\text{Se-}$, and $-\text{NR}_a$; a_4 and b_4 vary from 0 to 5; a , b , d , f , h , i , and j independently vary from 1-10; c , e , g , and k independently vary from 1-100; R_a ,

R_b , R_c , and R_d are defined in the same manner as Y_4 ; T is either H or a negative charge.

The present invention also relates to the novel compounds comprising cyanine dyes of general Formula 5




Formula 5

wherein R_{37} , R_{38} , R_{39} , R_{40} , R_{41} , R_{42} , R_{43} , R_{44} , R_{45} , Y_5 , and Z_5 are independently selected from the group consisting of -H, C1-C10 alkoxy, C1-C10 polyalkoxyalkyl, C1-C20 polyhydroxyalkyl, C5-C20 polyhydroxyaryl, glucose derivatives of R groups, saccharides, amino, C1-C10 aminoalkyl, cyano, nitro, halogen, hydrophilic peptides, arylpolysulfonates, C1-C10 alkyl, C1-C10 aryl, -SO₃T, -CO₂T, -OH, -(CH₂)_aSO₃T, -(CH₂)_aOSO₃T, -(CH₂)_aNHSO₃T, -(CH₂)_aCO₂(CH₂)_bSO₃T, -(CH₂)_aOCO(CH₂)_bSO₃T, -(CH₂)_aCONH(CH₂)_bSO₃T, -(CH₂)_aNHCO(CH₂)_bSO₃T, -(CH₂)_aNHCONH(CH₂)_bSO₃T, -(CH₂)_aNHCSNH(CH₂)_bSO₃T, -(CH₂)_aOCONH(CH₂)_bSO₃T, -(CH₂)_aPO₃HT, -(CH₂)_aPO₃T₂, -(CH₂)_aOPO₃HT, -(CH₂)_aOPO₃T₂, -(CH₂)_aNHPO₃HT, -(CH₂)_aNHPO₃T₂, -(CH₂)_aCO₂(CH₂)_bPO₃HT, -(CH₂)_aCO₂(CH₂)_bPO₃T₂, -(CH₂)_aOCO(CH₂)_bPO₃HT, -(CH₂)_aOCO(CH₂)_bPO₃T₂, -(CH₂)_aCONH(CH₂)_bPO₃HT, -(CH₂)_aCONH(CH₂)_bPO₃T₂, -(CH₂)_aNHCO(CH₂)_bPO₃HT, -(CH₂)_aNHCO(CH₂)_bPO₃T₂, -(CH₂)_aNHCONH(CH₂)_bPO₃HT,

- $-(CH_2)_aNHCONH(CH_2)_bPO_3T_2$, $-(CH_2)_aNHCSNH(CH_2)_bPO_3HT$,
 $-(CH_2)_aNHCSNH(CH_2)_bPO_3T_2$, $-(CH_2)_aOCONH(CH_2)_bPO_3HT$, and
 $-(CH_2)_aOCONH(CH_2)_bPO_3T_2$, $-CH_2(CH_2-O-CH_2)_c-CH_2-OH$, $-(CH_2)_d-CO_2T$,
 $-CH_2-(CH_2-O-CH_2)_e-CH_2-CO_2T$, $-(CH_2)_f-NH_2$, $-CH_2-(CH_2-O-CH_2)_g-CH_2-NH_2$,
5 $-(CH_2)_h-N(R_a)-(CH_2)_i-CO_2T$, and $-(CH_2)_j-N(R_b)-CH_2-(CH_2-O-CH_2)_k-CH_2-CO_2T$;
 W_5 and X_5 are selected from the group consisting of $-CR_cR_d$, $-O-$, $-NR_c$, $-S-$,
and $-Se$; V_5 is a single bond or is selected from the group consisting of $-O-$,
 $-S-$, $-Se-$, and $-NR_a$; D_5 is a single or a double bond; A_5 , B_5 and E_5 may be the
same or different and are selected from the group consisting of $-O-$, $-S-$, $-Se-$,
10 $-P-$, $-NR_a$, $-CR_cR_d$, CR_c , alkyl, and $-C=O$; A_5 , B_5 , D_5 , and E_5 may together form a
6 or 7 membered carbocyclic ring or a 6 or 7 membered heterocyclic ring
optionally containing one or more oxygen, nitrogen, or a sulfur atom; a , b , d , f ,
 h , i , and j independently vary from 1-10; c , e , g , and k independently vary from
1-100; a_5 and b_5 vary from 0 to 5; R_a , R_b , R_c , and R_d are defined in the same
15 manner as Y_5 ; T is either H or a negative charge.

The present invention also relates to the novel compounds comprising cyanine dyes of general Formula 6



Formula 6

wherein R_{46} , R_{47} , R_{48} , R_{49} , R_{50} , R_{51} , R_{52} , R_{53} , R_{54} , R_{55} , R_{56} , R_{57} and R_{58} , Y_6 , and Z_6 are independently selected from the group consisting of -H, C1-C10 alkoxyl, C1-C10 polyalkoxyalkyl, C1-C20 polyhydroxyalkyl, C5-C20 polyhydroxyaryl, glucose derivatives of R groups, saccharides, amino, C1-C10 aminoalkyl, cyano, nitro, halogen, hydrophilic peptides, arylpolysulfonates, C1-C10 alkyl, C1-C10 aryl, $-\text{SO}_3\text{T}$, $-\text{CO}_2\text{T}$, $-\text{OH}$, $-(\text{CH}_2)_a\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{OSO}_3\text{T}$, $-(\text{CH}_2)_a\text{NH}\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{CO}_2(\text{CH}_2)_b\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{OCO}(\text{CH}_2)_b\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{CONH}(\text{CH}_2)_b\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{NHCO}(\text{CH}_2)_b\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{NHCONH}(\text{CH}_2)_b\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{NHCSNH}(\text{CH}_2)_b\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{OCONH}(\text{CH}_2)_b\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{PO}_3\text{HT}$, $-(\text{CH}_2)_a\text{PO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{OPO}_3\text{HT}$, $-(\text{CH}_2)_a\text{OPO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{NHPO}_3\text{HT}$, $-(\text{CH}_2)_a\text{NHPO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{CO}_2(\text{CH}_2)_b\text{PO}_3\text{HT}$, $-(\text{CH}_2)_a\text{CO}_2(\text{CH}_2)_b\text{PO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{OCO}(\text{CH}_2)_b\text{PO}_3\text{HT}$, $-(\text{CH}_2)_a\text{OCO}(\text{CH}_2)_b\text{PO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{CONH}(\text{CH}_2)_b\text{PO}_3\text{HT}$, $-(\text{CH}_2)_a\text{CONH}(\text{CH}_2)_b\text{PO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{NHCO}(\text{CH}_2)_b\text{PO}_3\text{HT}$, $-(\text{CH}_2)_a\text{NHCO}(\text{CH}_2)_b\text{PO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{NHCONH}(\text{CH}_2)_b\text{PO}_3\text{HT}$, $-(\text{CH}_2)_a\text{NHCONH}(\text{CH}_2)_b\text{PO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{NHCSNH}(\text{CH}_2)_b\text{PO}_3\text{HT}$, $-(\text{CH}_2)_a\text{NHCSNH}(\text{CH}_2)_b\text{PO}_3\text{T}_2$, $-(\text{CH}_2)_a\text{OCONH}(\text{CH}_2)_b\text{PO}_3\text{HT}$, and $-(\text{CH}_2)_a\text{OCONH}(\text{CH}_2)_b\text{PO}_3\text{T}_2$, $-\text{CH}_2(\text{CH}_2\text{-O-CH}_2)_c\text{-CH}_2\text{-OH}$, $-(\text{CH}_2)_d\text{-CO}_2\text{T}$, $-\text{CH}_2\text{-(CH}_2\text{-O-CH}_2)_e\text{-CH}_2\text{-CO}_2\text{T}$, $-(\text{CH}_2)_f\text{-NH}_2$, $-\text{CH}_2\text{-(CH}_2\text{-O-CH}_2)_g\text{-CH}_2\text{-NH}_2$, $-(\text{CH}_2)_h\text{-N(R}_a\text{)-(CH}_2)_i\text{-CO}_2\text{T}$, and $-(\text{CH}_2)_j\text{-N(R}_b\text{)-CH}_2\text{-(CH}_2\text{-O-CH}_2)_k\text{-CH}_2\text{-CO}_2\text{T}$; W_6 and X_6 are selected from the group consisting of $-\text{CR}_c\text{R}_d$, $-\text{O-}$, $-\text{NR}_c$, $-\text{S-}$, and $-\text{Se-}$; V_6 is a single bond or is selected from the group consisting of $-\text{O-}$, $-\text{S-}$, $-\text{Se-}$, and $-\text{NR}_a$; D_6 is a single or a double bond; A_6 , B_6 and E_6 may be the same or different and are selected from the group consisting of $-\text{O-}$, $-\text{S-}$, $-\text{Se-}$, $-\text{P-}$, $-\text{NR}_a$, $-\text{CR}_c\text{R}_d$, CR_c , alkyl, and $-\text{C=O}$; A_6 , B_6 , D_6 , and E_6 may together form a

6 or 7 membered carbocyclic ring or a 6 or 7 membered heterocyclic ring optionally containing one or more oxygen, nitrogen, or sulfur atom; a, b, d, f, h, i, and j independently vary from 1-10; c, e, g, and k independently vary from 1-100; a₆ and b₆ vary from 0 to 5; R_a, R_b, R_c, and R_d are defined in the same manner as Y₆; T is either H or a negative charge.

A hydrophilic chelate such as ethylenediaminetetraacetic acid (EDTA), diethylenetriaminepentaacetic acid (DPTA), 1,4,7,10-tetraazacyclododecanetetraacetic acid (DOTA), or their derivatives, can be attached to the compounds of Formulas 1-6 as one or more R groups. These structures are expected to be highly water soluble.

The inventive compounds, compositions, and methods are advantageous since they provide a real-time, accurate, repeatable measure of renal excretion rate using exogenous markers under specific yet changing circumstances. This represents a substantial improvement over any currently available or widely practiced method, since currently, no reliable, continuous, repeatable bedside method for the assessment of specific renal function by optical methods exists. Moreover, since the inventive method depends solely on the renal elimination of the exogenous chemical entity, the measurement is absolute and requires no subjective interpretation based on age, muscle mass, blood pressure, etc. In fact it represents the nature of renal function in a particular patient, under these particular circumstances, at a precise moment in time.

The inventive compounds, compositions, and methods provide simple, efficient, and effective monitoring of organ function. The compound is administered and a sensor, either external or internal, is used to detect

absorption and/or emission to determine the rate at which the compound is cleared from the blood. By altering the R groups, the compounds may be rendered more organ specific.

BRIEF DESCRIPTION OF THE DRAWINGS

5 Figure 1: Reaction pathway for the preparation of indole derivatives.

 Figure 2: Reaction pathway for the preparation of benzindole derivatives.

 Figure 3: Reaction pathway for the preparation of
10 indocarbocyanine derivatives.

 Figure 4: Reaction pathway for the preparation of benzindocarbocyanine derivatives.

 Figure 5: Reaction pathway for the preparation of robust indocarbocyanine derivatives.

15 Figure 6: Reaction pathway for the preparation of robust benzindocarbocyanine derivatives.

 Figure 7: Reaction pathway for the preparation of long-wavelength absorbing indocarbocyanine derivatives.

 Figure 8A: Absorption spectrum of indoledisulfonate in water.

20 Figure 8B: Emission spectrum of indoledisulfonate in water.

 Figure 9A: Absorption spectrum of indocarbocyaninetetrasulfonate in water.

 Figure 9B: Emission spectrum of indocarbocyaninetetrasulfonate in water.

Figure 10A: Absorption spectrum of chloroindocarbocyanine in acetonitrile.

Figure 10B: Emission spectrum of chloroindocarbocyanine in acetonitrile.

5 Figure 11: Blood clearance profile of carbocyanine-polyaspartic (10 kDa) acid conjugate in a rat.

Figure 12: Blood clearance profile of carbocyanine-polyaspartic (30 kDa) acid conjugate in a rat.

Figure 13: Blood clearance profile of indoledisulfonate in a rat.

10 Figure 14: Blood clearance profile of carbocyaninetetrasulfonates in a rat.

DETAILED DESCRIPTION

In one embodiment the dyes of the invention serve as probes for continuous monitoring of renal function, especially for critically ill patients and
15 kidney transplant patients.

In another embodiment, the dyes of the invention are useful for dynamic hepatic function monitoring, especially for critically ill patients and liver transplant patients.

In another embodiment, the dyes of the invention are useful for
20 real-time determination of cardiac function, especially in patients with cardiac diseases.

In another embodiment, the dyes of the invention are useful for monitoring organ perfusion, especially for critically ill, cancer, and organ transplant patients.

In another embodiment, the dyes are useful for assessing the functional status of tumors and for monitoring tumor perfusion, such as in renal or hepatic cancer patients.

The novel dyes of the present invention are prepared according to the methods well known in the art, as illustrated in general in Figures 1-7 and described for specific compounds in Examples 1-11.

In one embodiment, the novel compounds, also called tracers, of the present invention have the Formula 1, wherein R_3 , R_4 , R_5 , R_6 and R_7 , and Y_1 are independently selected from the group consisting of -H, C1-C5 alkoxy, C1-C5 polyalkoxyalkyl, C1-C10 polyhydroxyalkyl, C5-C20 polyhydroxyaryl, mono- and disaccharides, nitro, hydrophilic peptides, arylpolysulfonates, C1-C5 alkyl, C1-C10 aryl, $-\text{SO}_3\text{T}$, $-\text{CO}_2\text{T}$, $-\text{OH}$, $-(\text{CH}_2)_a\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{OSO}_3\text{T}$, $-(\text{CH}_2)_a\text{NHSO}_3\text{T}$, $-(\text{CH}_2)_a\text{CO}_2(\text{CH}_2)_b\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{OCO}(\text{CH}_2)_b\text{SO}_3\text{T}$, $-\text{CH}_2(\text{CH}_2\text{-O-CH}_2)_c\text{-CH}_2\text{-OH}$, $-(\text{CH}_2)_d\text{-CO}_2\text{T}$, $-\text{CH}_2$ $-(\text{CH}_2\text{-O-CH}_2)_e\text{-CH}_2\text{-CO}_2\text{T}$, $-(\text{CH}_2)_f\text{-NH}_2$, $-\text{CH}_2\text{-(CH}_2\text{-O-CH}_2)_g\text{-CH}_2\text{-NH}_2$, $-(\text{CH}_2)_h$ $-\text{N(R}_a\text{)-(CH}_2)_i\text{-CO}_2\text{T}$, and $-(\text{CH}_2)_j\text{-N(R}_b\text{)-CH}_2\text{-(CH}_2\text{-O-CH}_2)_k\text{-CH}_2\text{-CO}_2\text{T}$; W_1 is selected from the group consisting of $-\text{CR}_c\text{R}_d$, $-\text{O-}$, $-\text{NR}_c$, $-\text{S-}$, and $-\text{Se-}$; a , b , d , f , h , i , and j independently vary from 1-5; c , e , g , and k independently vary from 1-20; R_a , R_b , R_c , and R_d are defined in the same manner as Y_1 ; T is a negative charge.

In another embodiment, the novel compounds of the present invention have the general Formula 2, wherein R_8 , R_9 , R_{10} , R_{11} , R_{12} , R_{13} , R_{14} , and Y_2 are independently selected from the group consisting of -H, C1-C5 alkoxy, C1-C5 polyalkoxyalkyl, C1-C10 polyhydroxyalkyl, C5-C20

polyhydroxyaryl, mono- and disaccharides, nitro, hydrophilic peptides,
arylpolysulfonates, C1-C5 alkyl, C1-C10 aryl, -SO₃T, -CO₂T, -OH,
-(CH₂)_aSO₃T, -(CH₂)_aOSO₃T, -(CH₂)_aNHSO₃T, -(CH₂)_aCO₂(CH₂)_bSO₃T,
-(CH₂)_aOCO(CH₂)_bSO₃T, -CH₂(CH₂-O-CH₂)_c-CH₂-OH, -(CH₂)_d-CO₂T, -CH₂
5 -(CH₂-O-CH₂)_e-CH₂-CO₂T, -(CH₂)_f-NH₂, -CH₂-(CH₂-O-CH₂)_g-CH₂-NH₂, -(CH₂)_h
-N(R_a)-(CH₂)_i-CO₂T, and -(CH₂)_j-N(R_b)-CH₂-(CH₂-O-CH₂)_k-CH₂-CO₂T; W₂ is
selected from the group consisting of -CR_cR_d, -O-, -NR_c, -S-, and -Se; a, b, d, f,
h, i, and j independently vary from 1-5; c, e, g, and k independently vary from 1-
20; R_a, R_b, R_c, and R_d are defined in the same manner as Y₂; T is a negative
10 charge.

In another embodiment, the novel compositions of the present
invention have the general Formula 3, wherein R₁₅, R₁₆, R₁₇, R₁₈, R₁₉, R₂₀, R₂₁,
R₂₂, R₂₃, Y₃, and Z₃ are independently selected from the group consisting of -H,
C1-C5 alkoxy, C1-C5 polyalkoxyalkyl, C1-C10 polyhydroxyalkyl, C5-C20
15 polyhydroxyaryl, mono- and disaccharides, nitro, hydrophilic peptides,
arylpolysulfonates, C1-C5 alkyl, C1-C10 aryl, -SO₃T, -CO₂T, -OH,
-(CH₂)_aSO₃T, -(CH₂)_aOSO₃T, -(CH₂)_aNHSO₃T, -(CH₂)_aCO₂(CH₂)_bSO₃T,
-(CH₂)_aOCO(CH₂)_bSO₃T, -CH₂(CH₂-O-CH₂)_c-CH₂-OH, -(CH₂)_d-CO₂T, -CH₂
-(CH₂-O-CH₂)_e-CH₂-CO₂T, -(CH₂)_f-NH₂, -CH₂-(CH₂-O-CH₂)_g-CH₂-NH₂, -(CH₂)_h
20 -N(R_a)-(CH₂)_i-CO₂T, and -(CH₂)_j-N(R_b)-CH₂-(CH₂-O-CH₂)_k-CH₂-CO₂T; W₃ and
X₃ are selected from the group consisting of -CR_cR_d, -O-, -NR_c, -S-, and -Se; V₃
is a single bond or is selected from the group consisting of -O-, -S-, -Se-, and
-NR_a; a, b, d, f, h, i, and j independently vary from 1-5; c, e, g, and k
independently vary from 1-50; a₃ and b₃ vary from 0 to 5; R_a, R_b, R_c, and R_d are
25 defined in the same manner as Y₃; T is either H or a negative charge.

In another embodiment, the novel compounds of the present invention have the general Formula 4, wherein R_{24} , R_{25} , R_{26} , R_{27} , R_{28} , R_{29} , R_{30} , R_{31} , R_{32} , R_{33} , R_{34} , R_{35} , R_{36} , Y_4 , and Z_4 are independently selected from the group consisting of -H, C1-C5 alkoxy, C1-C5 polyalkoxyalkyl, C1-C10 polyhydroxyalkyl, C5-C20 polyhydroxyaryl, mono- and disaccharides, nitro, hydrophilic peptides, arylpolysulfonates, C1-C5 alkyl, C1-C10 aryl, $-\text{SO}_3\text{T}$, $-\text{CO}_2\text{T}$, $-\text{OH}$, $-(\text{CH}_2)_a\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{OSO}_3\text{T}$, $-(\text{CH}_2)_a\text{NHSO}_3\text{T}$, $-(\text{CH}_2)_a\text{CO}_2(\text{CH}_2)_b\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{OCO}(\text{CH}_2)_b\text{SO}_3\text{T}$, $-\text{CH}_2(\text{CH}_2\text{-O-CH}_2)_c\text{-CH}_2\text{-OH}$, $-(\text{CH}_2)_d\text{-CO}_2\text{T}$, $-\text{CH}_2\text{-(CH}_2\text{-O-CH}_2)_e\text{-CH}_2\text{-CO}_2\text{T}$, $-(\text{CH}_2)_f\text{-NH}_2$, $-\text{CH}_2\text{-(CH}_2\text{-O-CH}_2)_g\text{-CH}_2\text{-NH}_2$, $-(\text{CH}_2)_h\text{-N(R}_a\text{)-(CH}_2)_i\text{-CO}_2\text{T}$, and $-(\text{CH}_2)_j\text{-N(R}_b\text{)-CH}_2\text{-(CH}_2\text{-O-CH}_2)_k\text{-CH}_2\text{-CO}_2\text{T}$; W_4 and X_4 are selected from the group consisting of $-\text{CR}_c\text{R}_d$, $-\text{O-}$, $-\text{NR}_c$, $-\text{S-}$, and $-\text{Se-}$; V_4 is a single bond or is selected from the group consisting of $-\text{O-}$, $-\text{S-}$, $-\text{Se-}$, and $-\text{NR}_a$; a_4 and b_4 vary from 0 to 5; a , b , d , f , h , i , and j independently vary from 1-5; c , e , g , and k independently vary from 1-50; R_a , R_b , R_c , and R_d are defined in the same manner as Y_4 ; T is either H or a negative charge.

In another embodiment, the novel compounds of the present invention have the general Formula 5, wherein R_{37} , R_{38} , R_{39} , R_{40} , R_{41} , R_{42} , R_{43} , R_{44} , R_{45} , Y_5 , and Z_5 are independently selected from the group consisting of -H, C1-C5 alkoxy, C1-C5 polyalkoxyalkyl, C1-C10 polyhydroxyalkyl, C5-C20 polyhydroxyaryl, mono- and disaccharides, nitro, hydrophilic peptides, arylpolysulfonates, C1-C5 alkyl, C1-C10 aryl, $-\text{SO}_3\text{T}$, $-\text{CO}_2\text{T}$, $-\text{OH}$, $-(\text{CH}_2)_a\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{OSO}_3\text{T}$, $-(\text{CH}_2)_a\text{NHSO}_3\text{T}$, $-(\text{CH}_2)_a\text{CO}_2(\text{CH}_2)_b\text{SO}_3\text{T}$, $-(\text{CH}_2)_a\text{OCO}(\text{CH}_2)_b\text{SO}_3\text{T}$, $-\text{CH}_2(\text{CH}_2\text{-O-CH}_2)_c\text{-CH}_2\text{-OH}$, $-(\text{CH}_2)_d\text{-CO}_2\text{T}$, $-\text{CH}_2\text{-(CH}_2\text{-O-CH}_2)_e\text{-CH}_2\text{-CO}_2\text{T}$, $-(\text{CH}_2)_f\text{-NH}_2$, $-\text{CH}_2\text{-(CH}_2\text{-O-CH}_2)_g\text{-CH}_2\text{-NH}_2$, $-(\text{CH}_2)_h$

-N(R_a)-(CH₂)_i-CO₂T, and -(CH₂)_j-N(R_b)-CH₂-(CH₂-O-CH₂)_k-CH₂-CO₂T; W₅ and X₅ are selected from the group consisting of -CR_cR_d, -O-, -NR_c, -S-, and -Se; V₅ is a single bond or is selected from the group consisting of -O-, -S-, -Se-, and -NR_a D₅ is a single or a double bond; A₅, B₅ and E₅ may be the same or
5 different and are selected from the group consisting of -O-, -S-, -NR_a, -CR_cR_d, CR_c, and alkyl; A₅, B₅, D₅, and E₅ may together form a 6 or 7 membered carbocyclic ring or a 6 or 7 membered heterocyclic ring optionally containing one or more oxygen, nitrogen, or sulfur atom; a, b, d, f, h, i, and j independently vary from 1-5; c, e, g, and k independently vary from 1-50; a₅ and b₅ vary from 0
10 to 5; R_a, R_b, R_c, and R_d are defined in the same manner as Y₅; T is either H or a negative charge.

In yet another embodiment, the novel compounds of the present invention have the general Formula 6, wherein R₄₆, R₄₇, R₄₈, R₄₉, R₅₀, R₅₁, R₅₂, R₅₃, R₅₄, R₅₅, R₅₆, R₅₇, R₅₈, Y₆, and Z₆ are independently selected from the
15 group consisting of -H, C1-C5 alkoxy, C1-C5 polyalkoxyalkyl, C1-C10 polyhydroxyalkyl, C5-C20 polyhydroxyaryl, mono- and disaccharides, nitro, hydrophilic peptides, arylpolysulfonates, C1-C5 alkyl, C1-C10 aryl, -SO₃T, -CO₂T, -OH, -(CH₂)_aSO₃T, -(CH₂)_aOSO₃T, -(CH₂)_aNHSO₃T, -(CH₂)_aCO₂(CH₂)_bSO₃T, -(CH₂)_aOCO(CH₂)_bSO₃T, -CH₂(CH₂-O-CH₂)_c-CH₂-OH,
20 -(CH₂)_d-CO₂T, -CH₂-(CH₂-O-CH₂)_e-CH₂-CO₂T, -(CH₂)_f-NH₂, -CH₂-(CH₂-O-CH₂)_g-CH₂-NH₂, -(CH₂)_h-N(R_a)-(CH₂)_i-CO₂T, and -(CH₂)_j-N(R_b)-CH₂-(CH₂-O-CH₂)_k-CH₂-CO₂T; W₆ and X₆ are selected from the group consisting of -CR_cR_d, -O-, -NR_c, -S-, and -Se; V₆ is a single bond or is selected from the group consisting of -O-, -S-, -Se-, and -NR_a; D₆ is a single or a double bond; A₆,
25 B₆ and E₆ may be the same or different and are selected from the group

consisting of -O-, -S-, -NR_a, -CR_cR_d, CR_c, and alkyl; A₆, B₆, D₆, and E₆ may together form a 6 or 7 membered carbocyclic ring or a 6 or 7 membered heterocyclic ring optionally containing one or more oxygen, nitrogen, or sulfur atom; a, b, d, f, h, i, and j independently vary from 1-5; c, e, g, and k
5 independently vary from 1-50; a₅ and b₅ vary from 0 to 5; R_a, R_b, R_c, and R_d are defined in the same manner as Y₆; T is either H or a negative charge.

The compounds of the invention can be formulated into diagnostic and therapeutic compositions for enteral or parenteral administration. These compositions contain an effective amount of the dye along with conventional
10 pharmaceutical carriers and excipients appropriate for the type of administration contemplated. For example, parenteral formulations advantageously contain the inventive agent in a sterile aqueous solution or suspension. Parenteral compositions may be injected directly or mixed with a large volume parenteral composition for systemic administration. Such
15 solutions also may contain pharmaceutically acceptable buffers and, optionally, electrolytes such as sodium chloride.

Formulations for enteral administration may vary widely, as is well known in the art. In general, such formulations are liquids, which include an effective amount of the inventive agent in aqueous solution or suspension.
20 Such enteral compositions may optionally include buffers, surfactants, thixotropic agents, and the like. Compositions for oral administration may also contain flavoring agents and other ingredients for enhancing their organoleptic qualities.

The compositions are administered in doses effective to achieve
25 the desired effect or result. The dosage of the tracers may vary according to

the clinical procedure contemplated and generally ranges from 1 picomolar to 100 millimolar. The compositions may be administered to a patient, typically a warm-blooded animal either systemically or locally to the organ or tissue to be imaged, and the patient then subject to the imaging procedure. The tracers
5 may be administered to the patient by any suitable method, including intravenous, intraperitoneal, or subcutaneous injection or infusion, oral administration, transdermal absorption through the skin, aerosols, or by inhalation. The detection of the tracers is achieved by optical fluorescence, absorbance, or light scattering methods known in the art (Muller et al. Eds.,
10 Medical Optical Tomography, SPIE Volume IS11, 1993, which is expressly incorporated herein by reference) using invasive or non-invasive probes such as endoscopes, catheters, ear clips, hand bands, surface coils, finger probes, and the like. Physiological function is correlated with the clearance profiles and rates of these agents from body fluids (R.B. Dorshow et al., Non-Invasive
15 Fluorescence Detection of Hepatic and Renal Function, *Bull. Am. Phys. Soc.* 1997, 42, 681, which is expressly incorporated by reference herein).

The inventive composition may be administered for imaging by more than one modality. As one example, the composition may be used for imaging by optical imaging alone, by nuclear imaging alone, or by both optical
20 and nuclear imaging modalities when a radioactive isotope is included in the chemical formula, such as replacing a halogen atom with a radioactive halogen, and/or including a radioactive metal ion such as Tc^{99} , In^{111} , etc. As another example, the composition may be used for imaging by optical imaging alone, by magnetic resonance (MR) alone, or by both optical and MR modalities when a

paramagnetic metal ion such as gadolinium or manganese is included in the chemical formula.

It will also be appreciated that the inventive compositions may be administered with other contrast agents or media used to enhance an image from a non-optical modality. These include agents for enhancing an image obtained by modalities including but not limited to MR, ultrasound (US), x-ray, positron emission tomography (PET), computed tomography (CT), single photon emission computed tomography (SPECT), optoacoustic (e.g. U.S. Patent Nos. 5,840,023 and 5,977,538 which are expressly incorporated by reference herein in their entirety), etc. Both optical and non-optical agents may be formulated as a single composition (that is, one composition containing one, two, or more components, for example, an optical agent and a MR agent), or may be formulated as separate compositions. The inventive optical imaging contrast agent and the non-optical contrast agent are administered in doses effective to achieve the desired enhancement, diagnosis, therapy, etc., as known to one skilled in the art. The inventive compositions, either alone or combined with a contrast agent, may be administered to a patient, typically a warm-blooded animal, systemically or locally to the organ or tissue to be imaged. The patient is then imaged by optical imaging and/or by another modality. As one example of this embodiment, the inventive compounds may be added to contrast media compositions. As another example, the inventive compositions may be co-administered with contrast media, either simultaneously or within the same diagnostic and/or therapeutic procedure (for example, administering the inventive composition and administering a contrast agent then performing optical imaging followed by another imaging modality, or

administering the inventive composition and administering a contrast agent then performing another imaging modality followed by optical imaging, or administering the inventive composition and optical imaging, then administering a contrast agent and MR, US, CT, etc. imaging, or administering a contrast agent and imaging by MR, US, CT, etc., then administering the inventive composition and optical imaging, or administering the inventive composition and a contrast agent, and simultaneously imaging by an optical modality and MR, US, CT, etc.). As another example, an optical imaging agent may be added as an additive or excipient for a non-optical imaging modality. In this embodiment, the optically active component, such as the dyes disclosed herein, could be added as a buffering agent to control pH or as a chelate to improve formulation stability, etc. in MR contrast media, CT contrast media, x-ray contrast media, US contrast media, etc. The MR, CT, x-ray, US contrast media would then also function as an optical imaging agent. The information obtained from the modality using the non-optical contrast agent is useful in combination with the image obtained using the optical contrast agent.

In one embodiment, the agents may be formulated as micelles, liposomes, microcapsules, or other microparticles. These formulations may enhance delivery, and localization of the inventive compounds to/at the desired organ or site. The target specificity of these formulations can be enhanced by using suitable targeting molecules such as peptides, saccharides, fatty acids, etc. Preparation and loading of these are well known in the art.

As one example, liposomes may be prepared from dipalmitoyl phosphatidylcholine (DPPC) or egg phosphatidylcholine (PC) because this lipid has a low heat transition. Liposomes are made using standard procedures as

known to one skilled in the art (e.g., Braun-Falco et al., (Eds.), Griesbach Conference, Liposome Dermatics, Springer-Verlag, Berlin (1992)).

Polycaprolactone, poly(glycolic) acid, poly(lactic) acid, polyanhydride or lipids may be formulated as microspheres. As an illustrative example, the optical

5 agent may be mixed with polyvinyl alcohol (PVA), the mixture then dried and coated with ethylene vinyl acetate, then cooled again with PVA. In a liposome, the optical agent may be within one or both lipid bilayers, in the aqueous between the bilayers, or with the center or core. Liposomes may be modified with other molecules and lipids to form a cationic liposome. Liposomes may
10 also be modified with lipids to render their surface more hydrophilic which increases their circulation time in the bloodstream. The thus-modified liposome has been termed a "stealth" liposome, or a long-lived liposome, as described in U.S. Patent Nos. 6,277,403; 6,610,322; 5,631,018; 5,395,619; and 6,258,378, each of which is expressly incorporated by reference herein in its entirety, and
15 in Stealth Liposomes, Lasic and Martin (Eds.) 1995, CRC Press, London.

Encapsulation methods include detergent dialysis, freeze drying, film forming, injection, as known to one skilled in the art and disclosed in, for example, U.S. Patent No. 6,406,713 which is expressly incorporated by reference herein in its entirety.

20 The agent formulated in liposomes, microcapsules, etc. may be administered by any of the routes previously described. In a formulation applied topically, the optical agent is slowly released over time. In an injectable formulation, the liposome capsule circulates in the bloodstream and is delivered to a desired site.

As another example, microparticles such as ultra small iron oxide particles (USPIO) and other metallic particles such as silver or gold particles coated with or attached (covalently or non-covalently) with the inventive compounds may be used for optical imaging and/or MRI. Such particles are
5 known to one skilled in the art as disclosed in, for example, U.S. Patent No. 5,492,814 and Journal of Biomedical Optics 8(3), 472-478 (July 2003) which are expressly incorporated by reference herein in their entirety.

Organ function can be assessed either by the differences in the manner in which the normal and impaired cells remove the tracer from the
10 bloodstream, by measuring the rate or accumulation of these tracers in the organs or tissues, or by obtaining tomographic images of the organs or tissues. Blood pool clearance may be measured non-invasively from convenient surface capillaries such as those found in an ear lobe or a finger, for example, using an ear clip or finger clip sensor, or may be measured invasively using an
15 endovascular catheter. Accumulation of the tracer within the cells of interest can be assessed in a similar fashion. The clearance of the tracer dyes may be determined by selecting excitation wavelengths and filters for the emitted photons. The concentration-time curves may be analyzed in real time by a microprocessor. In order to demonstrate feasibility of the inventive compounds
20 to monitor organ function, a non-invasive absorbance or fluorescence detection system to monitor the signal emanating from the vasculature infused with the compounds is used. Indole derivatives, such as those of Formulas 1-6, fluoresce at a wavelength between 350 nm and 1300 nm when excited at the appropriate wavelength as is known to, or readily determined by, one skilled in
25 the art.

In addition to the noninvasive techniques, a modified pulmonary artery catheter can be used to make the necessary measurements (R.B.

Dorshow, J.E. Bugaj, S.A. Achilefu, R. Rajagopalan, and A.H. Combs,

Monitoring Physiological Function by Detection of Exogenous Fluorescent

5 Contrast Agents, in *Optical Diagnostics of Biological Fluids IV*, A. Priezzhev and T. Asakura, Editors, Proceedings of SPIE 1999, 3599, 2-8, which is expressly incorporated by reference herein). Currently, pulmonary artery catheters measure only intravascular pressures, cardiac output and other derived measures of blood flow. Critically ill patients are managed using these
10 parameters, but rely on intermittent blood sampling and testing for assessment of renal function. These laboratory parameters represent discontinuous data and are frequently misleading in many patient populations. Yet, importantly, they are relied upon heavily for patient assessment, treatment decisions, and drug dosing.

15 The modified pulmonary artery catheter incorporates an optical sensor into the tip of a standard pulmonary artery catheter. This wavelength specific optical sensor can monitor the renal function specific elimination of an optically detectable chemical entity. Thus, by a method analogous to a dye dilution curve, real-time renal function can be monitored by the disappearance
20 of the optically detected compound. Modification of a standard pulmonary artery catheter only requires making the fiber optic sensor wavelength specific, as is known to one skilled in this art. Catheters that incorporate fiber optic technology for measuring mixed venous oxygen saturation currently exist.

The present invention may be used for rapid bedside evaluation
25 of renal function and also to monitor the efficiency of hemodialysis. The

invention is further demonstrated by the following examples. Since many modifications, variations, and changes in detail may be made to the described embodiments, it is intended that all matter in the foregoing description and shown in the accompanying drawings be interpreted as illustrative and not in a limiting sense.

EXAMPLE 1

Synthesis of indole disulfonate (Figure 1, Compound 5, $Y_7 = \text{SO}_3^-$; $X_7 = \text{H}$; $n = 1$)


10

A mixture of 3-methyl-2-butanone (25.2 mL), and p-hydrazinobenzenesulfonic acid (15 g) in acetic acid (45 mL) was heated at 110°C for 3 hours. After reaction, the mixture was allowed to cool to room temperature and ethyl acetate (100 mL) was added to precipitate the product, which was filtered and washed with ethyl acetate (100 mL). The intermediate compound, 2,3,3-trimethylindolenium-5-sulfonate (Figure 1, compound 3) was obtained as a pink powder in 80% yield. A portion of compound 3 (9.2 g) in methanol (115 mL) was carefully added to a solution of KOH in isopropanol (100 mL). A yellow potassium salt of the sulfonate was obtained in 85% yield after vacuum-drying for 12 hours. A portion of the 2,3,3-trimethylindolenium-5-sulfonate potassium salt (4 g) and 1,3-propanesultone (2.1 g) was heated in dichlorobenzene (40 mL) at 110°C for 12 hours. The mixture was allowed to cool to room temperature and the resulting precipitate was filtered and washed with isopropanol. The resulting pink powder was dried under vacuum to give 97% of the desired compound.


25

Other compounds prepared by a similar method described above

include polyhydroxyl indoles such as



and



5

EXAMPLE 2

Synthesis of indole disulfonate

(Figure 1, Compound 5, $Y_7 = \text{SO}_3^-$; $X_7 = \text{H}$; $n = 2$)

This compound was prepared by the same procedure described in Example 1,


10 except that 1,4-butanedisulfone was used in place of 1,3-propanedisulfone.

- 32 -
EXAMPLE 3


Synthesis of benzoindole disulfonate
(Figure 2, Compound 8, $Y_7, Y_8 = \text{SO}_3^-$; $X_7 = \text{H}$; $n = 2$)

5 This compound was prepared by the same procedure described in Example 1, except that hydrazinonaphthalenedisulfonic acid was used in place of hydrazinobenzenesulfonic acid.

Other compounds prepared by a similar method include polyhydroxyindoles such as:



and




EXAMPLE 4

Synthesis of benzoindole disulfonate
(Figure 2, Compound 8, $Y_7, Y_8 = \text{SO}_3^-$; $X_7 = \text{OH}$; $n = 4$)

5 This compound was prepared by the same procedure described in Example 1, except that 3-hydroxymethyl-4-hydroxyl-2-butanone was used in place of 3-methyl-2-butanone.

EXAMPLE 5

Synthesis of Bis(ethylcarboxymethyl)indocyanine Dye



A mixture of 1,1,2-trimethyl-[1H]-benz[e]indole (9.1 g, 43.58 mmol) and 3-bromopropanoic acid (10.0 g, 65.37 mmol) in 1,2-dichlorobenzene (40 mL) was heated at 110 °C for 12 hours. The solution was cooled to room temperature and the red residue obtained was filtered and washed with acetonitrile: diethyl ether (1:1) mixture. The solid obtained was dried under vacuum to give 10 g (64%) of light brown powder. A portion of this solid (6.0 g; 16.56 mmol), glutaconaldehyde dianil monohydrochloride (2.36 g, 8.28 mmol) and sodium acetate trihydrate (2.93 g, 21.53 mmol) in ethanol (150 mL) were refluxed for 90 minutes. After evaporating the solvent, 40 mL of 2 N aqueous HCl was added to the residue and the mixture was centrifuged and the supernatant was decanted. This procedure was repeated

15


20

until the supernatant became nearly colorless. About 5 mL of water:acetonitrile (3:2) mixture was added to the solid residue and lyophilized to obtain 2 g of dark green flakes. The purity of the compound was established with $^1\text{H-NMR}$ and liquid chromatography/mass spectrometry (LC/MS).

5

EXAMPLE 6

Synthesis of Bis(pentylcarboxymethyl)indocyanine Dye



A mixture of 2,2,3-trimethyl-[1H]-benz[e]indole (20 g, 95.6
10 mmoles) and 6-bromohexanoic acid (28.1 g, 144.1 mmoles) in 1,2-dichlorobenzene (250 mL) was heated at 110 C for 12 hours. The green solution was cooled to room temperature and the brown solid precipitate formed was collected by filtration. After washing the solid with 1,2-dichlorobenzene and diethyl ether, the brown powder obtained (24 g, 64%) was
15 dried under vacuum at room temperature. A portion of this solid (4.0 g; 9.8 mmoles), glutaconaldehyde dianil monohydrochloride (1.4 g, 5 mmoles) and sodium acetate trihydrate (1.8 g, 12.9 mmoles) in ethanol (80 mL) were

refluxed for 1 hour. After evaporating the solvent, 20 mL of a 2 N aqueous HCl was added to the residue and the mixture was centrifuged and the supernatant was decanted. This procedure was repeated until the supernatant became nearly colorless. About 5 mL of water:acetonitrile (3:2) mixture was added to the solid residue and lyophilized to obtain about 2 g of dark green flakes. The purity of the compound was established with $^1\text{H-NMR}$, HPLC, and LC-MS.

EXAMPLE 7

Synthesis of polyhydroxyindole sulfonate (Figure 3, Compound 13; $\text{Y}_7, \text{Y}_8 = \text{O}_3^-$; $\text{X}_7 = \text{OH}$; $n=2$)

10

Phosphorus oxychloride (37 ml, 0.4 mole) was added dropwise with stirring to a cooled (-2°C) mixture of dimethylformamide (DMF, 0.5 mole, 40 mL) and dichloromethane (DCM, 40 mL), followed by the addition of acetone (5.8 g, 0.1 mole). The ice bath was removed and the solution refluxed for 3 hours. After cooling to room temperature, the product was either partitioned in water/DCM, separated and dried, or was purified by fractional distillation. Nuclear magnetic resonance and mass spectral analyses showed that the desired intermediate, 10, was obtained. Reaction of the intermediate with 2 equivalents of 2,2,3-trimethyl-[H]-benz[e]indolesulfonate-N-propanoic acid and 2 equivalents of sodium acetate trihydrate in ethanol gave a blue-green solution after 1.5 hours at reflux. Further functionalization of the dye with bis(isopropylidene)acetal protected monosaccharide is effected by the method described in the literature (J. H. Flanagan, C. V. Owens, S. E. Romero, et al., Near infrared heavy-atom-modified fluorescent dyes for base-calling in DNA-sequencing application using temporal discrimination. *Anal. Chem.*, 1998, 70(13), 2676-2684).

25

- 36 -
EXAMPLE 8

Synthesis of polyhydroxyindole sulfonate
(Figure 4, Compound 16, $Y_7, Y_8 = SO_3^-$; $X_7 = H$; $n = 1$)

5 Preparation of this compound was readily accomplished by the same procedure described in Example 6 using p-hydroxybenzenesulfonic acid in the place of the monosaccharide, and benzoindole instead of indole derivatives.

10 EXAMPLE 9

Synthesis of polyhydroxyindole sulfonate
(Figure 5, Compound 20, $Y_7, Y_8 = H$; $X_7 = OH$; $n = 1$)

The hydroxyindole compound was readily prepared by a literature
15 method (P.L. Southwick, J.G. Cairns, L.A. Ernst, and A.S. Waggoner, One pot Fischer synthesis of (2,3,3-trimethyl-3-H-indol-5-yl)-acetic acid derivatives as intermediates for fluorescent biolabels, *Org. Prep. Proced. Int. Briefs*, 1988, 20(3), 279-284). Reaction of p-carboxymethylphenylhydrazine hydrochloride (30 mmol, 1 equiv.) and 1,1-bis(hydroxymethyl)propanone (45 mmol, 1.5
20 equiv.) in acetic acid (50 mL) at room temperature for 30 minutes and at reflux for 1 gave (3,3-dihydroxymethyl-2-methyl-3-H-indol-5-yl)-acetic acid as a solid residue.

The intermediate 2-chloro-1-formyl-3-hydroxymethylenecyclohexane was prepared as described in the literature (G. A. Reynolds and K. H.
25 Drexhage, Stable heptamethine pyrylium dyes that absorb in the infrared. *J. Org. Chem.*, 1977, 42(5), 885-888). Equal volumes (40 mL each) of dimethylformamide (DMF) and dichloromethane were mixed and the solution was cooled to $-10^{\circ}C$ in acetone-dry ice bath. Under argon atmosphere,

phosphorus oxychloride (40 mL) in dichloromethane was added dropwise to the cool DMF solution, followed by the addition of 10 g of cyclohexanone. The resulting solution was allowed to warm up to room temperature and heated at reflux for 6 hours. After cooling to room temperature, the mixture was poured
5 into ice-cold water and stored at 4°C for 12 hours. A yellow powder was obtained. Condensation of a portion of this cyclic dialdehyde (1 equivalent) with the indole intermediate (2 equivalents) was carried out as described in Example 5. Further, the functionalization of the dye with bis (isopropylidene)acetal protected monosaccharide was effected by the method described in the
10 literature (J. H. Flanagan, C. V. Owens, S. E. Romero, et al., Near infrared heavy-atom-modified fluorescent dyes for base-calling in DNA-sequencing application using temporal discrimination. *Anal. Chem.*, 1998, 70(13), 2676-2684).

15

EXAMPLE 10

Synthesis of polyhydroxylbenzoindole sulfonate (Figure 6, Compound 22, Y₇, Y₈=H; X₇=OH; n=1)

A similar method described in Example 8 was used to prepare this
20 compound by replacing the indole with benzoindole derivatives.

EXAMPLE 11

Synthesis of rigid heteroatomic indole sulfonate (Figure 7, Compound 27, Y₇, Y₈, X₇= H; n = 1)

25

Starting with 3-oxo-4-cyclohexenone, this heteroatomic hydrophilic dye was readily prepared as described in Example 8.

- 38 -
EXAMPLE 12

Minimally invasive monitoring of the blood clearance profile of the dyes

A laser of appropriate wavelength for excitation of the dye chromophore was directed into one end of a fiber optic bundle and the other
5 end was positioned a few millimeters from the ear of a rat. A second fiber optic bundle was also positioned near the same ear to detect the emitted fluorescent light, and the other end was directed into the optics and electronics for data collection. An interference filter (IF) in the collection optics train was used to select emitted fluorescent light of the appropriate wavelength for the dye
10 chromophore.

Sprague-Dawley or Fischer 344 rats were anesthetized with urethane administered via intraperitoneal injection at a dose of 1.35 g/kg body weight. After the animals had achieved the desired plane of anesthesia, a 21 gauge butterfly with 12" tubing was placed in the lateral tail vein of each animal
15 and flushed with heparinized saline. The animals were placed onto a heating pad and kept warm throughout the entire study. The lobe of the left ear was affixed to a glass microscope slide to reduce movement and vibration.

Incident laser light delivered from the fiber optic was centered on the affixed ear. Data acquisition was then initiated, and a background reading
20 of fluorescence was obtained prior to administration of the test agent.

The compound was administered to the animal through a bolus injection in the lateral tail vein. The dose was typically 0.05 to 20 μ mole/kg of body weight. The fluorescence signal rapidly increased to a peak value, then decayed as a function of time as the conjugate cleared from the bloodstream.

This procedure was repeated with several dye-peptide conjugates in normal and tumored rats. Representative profiles are shown in Figures 6-10.

While the invention has been disclosed by reference to the details of preferred embodiments of the invention, it is to be understood that the disclosure is intended in an illustrative rather than in a limiting sense, as it is contemplated that modifications will readily occur to those skilled in the art, within the spirit of the invention and the scope of the appended claims.

What is claimed is: